

Ventilation/Perfusion Mismatch in Pulmonary Vein Stenosis Secondary to Atrial Fibrillation Ablation

Leah Anne Christine L. Bollos¹, Ryosuke Kasai², Hideki Otsuka^{2*}, Yoichi Otomi³, Tomomi Matsuura⁴, Tamaki Otani⁵, Koji Yamaguchi⁴, Takanori Bando¹, Yuya Ueki¹, Noritake Matsuda¹, Satoru Takashi³, Shota Azane³, Yamato Kunikane³, Shoichiro Takao⁶, Shusuke Yagi⁴, Masataka Sata⁴, Hitoshi Ikushima⁷, Masafumi Harada³

¹Tokushima University Graduate School of Health Sciences, Tokushima City, Japan

²Department of Medical Imaging/Nuclear Medicine, Tokushima University Graduate School of Biomedical Sciences, Tokushima City, Japan

³Department of Radiology, Tokushima University Hospital, Tokushima City, Japan

⁴Department of Cardiovascular Medicine, Tokushima University Hospital, Tokushima City, Japan

⁵Advance Radiation Research, Education and Management Center, Tokushima University, Tokushima City, Japan

⁶Department of Diagnostic Radiology, Tokushima University Graduate School of Biomedical Sciences, Tokushima City, Japan

⁷Department of Radiation Oncology, Tokushima University Graduate School of Biomedical Sciences, Tokushima City, Japan

ARTICLE INFO

Article type:
Case Series

Article history:
Received: 9 May 2024
Revised: 2 July 2024
Accepted: 1 Aug 2024

Keywords:
Ventilation/ Perfusion Scan
Ventilation/ Perfusion Mismatch
Pulmonary Vein Stenosis
Atrial Fibrillation
Pulmonary Vein Ablation

ABSTRACT

We present two patients with a history of paroxysmal atrial fibrillation who developed pulmonary vein stenosis (PVS) following atrial fibrillation (AF) ablation. Case 1 involved a female patient in her 50s who was asymptomatic for pulmonary symptoms but was found to have a high degree of left superior PVS 15 months after AF ablation. This was demonstrated using contrast-enhanced computed tomography (CE-CT) and supported by findings of perfusion defects on ventilation-perfusion (V/Q) scan. Case 2 was a male patient in his 60s who developed progressive left superior PVS nine months after AF ablation, evidenced by serial CE-CT and V/Q scans.

PVS is a rare but well-known complication of pulmonary vein ablation for the treatment of AF that can lead to severe complications if left untreated. V/Q scans effectively assess the functional significance of PVS by detecting abnormal blood flow segments. Although a V/Q mismatch characterized by reduced perfusion defects is more commonly used in evaluating pulmonary embolism, PVS should not be disregarded as a differential diagnosis. Few studies emphasize the utility of V/Q scans in managing PVS and highlight V/Q mismatch as a notable finding. This case report aimed to highlight their significance.

► Please cite this paper as:

Bollos LACL, Kasai R, Otsuka H, Otomi Y, Matsuura T, Otani T, Yamaguchi K, Bando T, Ueki Y, Matsuda N, Takashi S, Azane Sh, Kunikane Y, Takao Sh, Yagi Sh, Sata M, Ikushima H, Harada M. Ventilation/Perfusion Mismatch in Pulmonary Vein Stenosis Secondary to Atrial Fibrillation Ablation. *Asia Ocean J Nucl Med Biol.* 2025; 13(1): 62-69. doi: 10.22038/aojnmb.2024.79650.1561

Introduction

Pulmonary vein stenosis (PVS) is a well-known complication of pulmonary vein ablation for treating atrial fibrillation (1). It typically manifests between 3 and 6 months post-procedure (1, 2) but can also occur later, depending on the stenosis severity and vessel

involvement (2–4). PVS arises from tissue injury caused by thermal application near the pulmonary veins, resulting in luminal narrowing due to myocardial fibrosis and vessel wall thickening(5).

Increased lobar wedge pressure in patients with >50% stenosis may cause non-specific

* Corresponding author: Hideki Otsuka. Kuramoto-cho 3-18-15, Tokushima City, Tokushima, Japan #770-8503, Japan. Tel: 81-88-633-7173; Email: hideki.otsuka@tokushima-u.ac.jp

© 2025 *mums.ac.ir* All rights reserved.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

symptoms such as exertional dyspnea, pleuritic chest pain, cough, hemoptysis, and, in advanced stages, pulmonary hypertension (1, 6). Due to improved ablation techniques and awareness of complication risk factors, the incidence of PVS has since decreased. Severe PVS (>65% luminal narrowing) is rare, occurring in 0-3.1% of cases (4, 7). In comparison, mild to moderate cases (<50% luminal narrowing) are observed in up to 31% of patients who undergo pulmonary vein isolation (PVI) (8). Hemodynamic compensation may mask symptoms in mild-to-moderate cases (9), causing the disease to be overlooked and the true incidence of PVS to be underestimated.

The consensus for managing severe symptomatic PVS secondary to atrial fibrillation ablation is percutaneous intervention using pulmonary vein angioplasty with or without stent placement (10). Opinions vary in managing asymptomatic cases, with some studies suggesting the clinical benefits of early intervention (5, 10). Delayed diagnosis may lead to life-threatening progression (11), including possible progressive pulmonary vein hypoplasia proximal to the stenosis and total occlusion, which may then render percutaneous intervention ineffective (2,3). Therefore, a high index of suspicion and proper imaging screening for at-risk patients are necessary (11).

Imaging in PVS

Findings on imaging modalities may be non-specific for PVS and often require anatomical and functional imaging for proper assessment. Noninvasive imaging modalities, including computed tomography (CT), magnetic resonance imaging (MRI), V/Q scan, and transesophageal echocardiography (TEE), are typically used for this purpose (1). CT angiography with a pulmonary vein protocol is considered well-suited for delineating pulmonary vein anatomy, diagnosing PVS, and planning procedures; however, it may overestimate PV occlusions and raise concerns about radiation and contrast doses, especially for serial imaging (7).

V/Q scans are effective for assessing the functional significance of PVS (6, 12). It can be performed using planar scintigraphy, single-photon emission computed tomography (SPECT), or SPECT combined with low-dose CT (SPECT/CT). V/Q scans are highly sensitive in detecting regions with abnormal blood flow by taking advantage of the unique pulmonary arterial segmental anatomy. Under normal conditions, the apparent blood flow in one-third of the lungs ranges from 15% to 30%. With severe PVS (>65% luminal narrowing), local

perfusion within the affected pulmonary vein can substantially decrease to levels as low as 3% to 4% (13, 14). The reduced blood flow is demonstrated using radionuclide perfusion (Q) scans, widely used to evaluate relative lung perfusion among pathologies with vascular occlusions such as pulmonary embolism, neoplastic processes, or pulmonary congenital abnormalities (6).

During the perfusion scan (Q), the radiotracer ^{99m}Tc-macroaggregated albumin (MAA) is intravenously injected and travels through the right atrium, right ventricle, and pulmonary artery, accumulating through micro-embolization in the peripheral pulmonary vascular bed. Significant occlusions lead to decreased distribution of ^{99m}Tc-MAA in the affected lung areas (6).

V/Q mismatch characterized by reduced perfusion defects with no corresponding ventilatory abnormality is more commonly used to evaluate the likelihood of pulmonary embolism (12). In the case of PVS, V/Q mismatch can help determine the progression and functional significance of the occluded vessel in conjunction with CT or MRI findings (6, 15). Few studies emphasize the utility of V/Q scans in managing PVS or highlight V/Q mismatch as a notable finding (4, 6, 15, 16). We present two cases with such findings and highlight the role of the V/Q scan in managing patients with PVS.

Case Reports

Case 1

A female patient in her 50s with a history of paroxysmal atrial fibrillation and previous sessions of atrial fibrillation ablation returned 15 months after a recent ablation procedure with complaints of palpitations. Despite being on antiarrhythmic and calcium channel blocker medications, she had prolonged atrial fibrillation episodes, as observed on Holter monitoring. She denied shortness of breath, exertional dyspnea, hemoptysis, or chest pain throughout this time. The patient was admitted for further investigation and possible atrial fibrillation ablation.

CE-CT and V/Q scans assessed morphological and functional abnormalities, respectively. Initial CT with 3D reconstruction suggested a high degree of stenosis at the LSPV inlet (Figure 1). A V/Q scan revealed normal bilateral ventilation (V) but with moderately decreased perfusion (Q) of the left upper lung (Figure 2). This V/Q mismatch defect (Figure 3) supports the CT scan findings of severe stenosis of the LSPV.

ICE-guided percutaneous transluminal angioplasty (PTA) was performed to dilate the

LSPV. Post-procedural imaging was conducted using a hybrid SPECT/CT scanner, where lung scintigraphy was performed (Figure 4). This revealed an improved perfusion in the left upper lobe, although a mild perfusion defect was still noted. The bilateral ventilation

remained normal. Consequently, the V/Q mismatch decreased; however, this was not entirely resolved (Figure 3). CT 3-D reconstruction imaging demonstrated the reestablishment of the LSPV (Figure 1).




Figure1. Case 1. 3-D Rendering of Pre- and Post-PTA CT Scan Images
Pre-PTA CT image shows an abrupt tapering and cut-off at the area of the left superior pulmonary vein (**outlined**), suggesting a filling defect from an occlusion. The post-PTA CT image shows the re-establishment of the LSPV (**outlined**)
Abbreviation: PTA- Percutaneous Transluminal Angioplasty




Figure2. Case 1. V/Q scan, Pre- and Post- PTA to LSPV
Pre-PTA V/Q scan revealed a moderately decreased perfusion (Q) of the left upper lung (**black arrows**) and normal bilateral ventilation (V). **White arrows** on the ventilation scan point toward the same area of the left upper lung with the perfusion defect. In comparison, post-PTA V/Q scan shows improved perfusion of the left upper lung with some mildly decreased perfused areas. Ventilation scan remain normal in both lungs. (a) Anterior view, (b) posterior view, (c) right lateral view, (d) left lateral view
Abbreviations: PTA- Percutaneous Transluminal Angioplasty, LSPV- Left Superior Pulmonary Vein




Figure 3. Case 1. V/Q mismatch image

Planar images show a posterior view of the patient’s perfusion scan (a) and ventilation scan (b), along with the respective representation of their mismatched areas, which are darkened (c). A comparison with images taken post PTA shows that the mismatched areas have diminished (arrows), although some areas remain unresolved. This is achieved by subtracting the pre-PTA perfusion image (d), from the post PTA perfusion image (e) or (post-PTA – pre-PTA). The perfusion subtracted image illustrates the resolved mismatched areas (f) (arrows)

Abbreviation: PTA- Percutaneous Transluminal Angioplasty




Figure 4. Case 1. Perfusion planar and MAA-SPECT/CT

The hybrid SPECT/ CT scanner enables both morphological and functional assessment of the lung. Morphological findings such as consolidation, ground glass opacity and reticular / fibrotic changes can be seen, while functional assessment using fused SPECT/CT images from any directions (axial, coronal, sagittal) provides better anatomic visualization at a higher resolution. In comparison with the planar images (a) & (b) are SPECT/CT (c) & (d) show more anatomical visualization of the lungs

Abbreviation: PTA- Percutaneous Transluminal Angioplasty

Case 2

A male patient in his 60s with a history of recurrent paroxysmal atrial fibrillation was found to have a PVS in the left superior pulmonary vein. This was observed on a CT scan 9 months after the second ablation procedure (Figure 5). Subsequent imaging conducted 2 and 3 months later revealed a ventilation-perfusion mismatch in the left upper lobe of the lung on the V/Q scan and a worsening of the

stenosis on the CE-CT scan (Figure 6). The patient did not present with any related symptoms despite imaging findings. PTA was then performed, and serial postprocedural imaging revealed a gradual improvement of lung perfusion on the affected lobe (Figure 7). On CT scan, an increase in vessel diameter and increased contrast enhancement of the left superior pulmonary vein were observed (17).




Figure 5. Case 2. Pre- and Post-Ablation CE-CT images

CE-CT in axial, sagittal and coronal views, taken 2 months post-ablation shows stenosis near the ostium of the LSPV (arrows). Comparison with pre-ablation images suggest the stenosis had developed after the procedure

Abbreviations: PTA- Percutaneous Transluminal Angioplasty, PVS- Pulmonary Vein Stenosis, CE-CT- Contrast Enhanced Computed Tomography, LSPV- Left Superior Pulmonary Vein




Figure 6. Case 2. Pre- Angioplasty Imaging

Perfusion (a) and ventilation (b) planar images demonstrate V/Q mismatch; Perfusion defects (a) are seen in the left upper lobe of the lung (black arrows) while corresponding areas in the ventilation scan (b) are preserved. V/Q mismatch (c) highlights the discrepancy of areas between the posterior and ventilation scan. Contrast-enhanced CT of the chest (d) in axial, sagittal and coronal views, reveal the LSPV tapering- off with minimal contrast enhancement proximal to the stenosed area noted (yellow arrows)

Abbreviations: PTA- Percutaneous Transluminal Angioplasty, AF- Atrial Fibrillation, PVS- Pulmonary Vein Stenosis, LSPV- Left Superior Pulmonary Vein, V/Q – Ventilation/ Perfusion, Ant. – Anterior View, L – Left, P – Posterior View, LL- Left Lateral View, RL- Right Lateral View




Figure 7. Case 2. Post- Angioplasty Images

Perfusion (a) and ventilation (b) planar images show V/Q mismatch has diminished; Perfusion of the left upper lobe of the lung (a) has partially recovered while corresponding areas in the ventilation scan (b) remain unchanged. The mismatched areas(c) have also reduced. Contrast-enhanced CT of the chest (d) in axial, sagittal and coronal views, reveal a reduction of the stenosis with partially reestablished connections that gradually improve further on 1 year follow up (e)

Abbreviations: PTA- Percutaneous Transluminal Angioplasty, AF- Atrial Fibrillation, PVS- Pulmonary Vein Stenosis, LSPV- Left Superior Pulmonary Vein, V/Q – Ventilation/ Perfusion, Ant. – Anterior View, L – Left, P – Posterior View, LL- Left Lateral View, RL- Right Lateral View

Discussion

We present two cases of PVS secondary to atrial fibrillation ablation in which CE-CT and V/Q scans were utilized for the morphological and functional assessment of PVS. Both cases exhibited a V/Q mismatch due to decreased perfusion in the upper left lung, coinciding with the CE-CT image findings.

When significantly occluded, PVS results in a stagnation of blood flow and decreased perfusion to the part of the lung supplied by the affected vessel. This consequently depicts mismatch defects due to decreased perfusion in the areas drained by the occluded vessel. This finding is not specific to PVS and can be found in conditions with pulmonary vessel obstruction, such as pulmonary embolism, malignancies, pulmonary hypertension, and COPD-related pulmonary hypertension (9, 18). Though V/Q scans may lack specificity for PVS, some studies have noted that patients with severe PVS (>70% luminal narrowing) consistently demonstrate decreased lung perfusion and are associated with poor clinical outcomes. It is, however, less noted in those with mild stenosis (less than 50%) (3, 4). Nevertheless, when V/Q scan findings are used in the proper clinical context, PVS should be considered as a differential diagnosis. V/Q scan can also be applied quantitatively and is frequently used to assess pulmonary perfusion defects and endovascular treatment response (19). They can also aid in determining the significance of an ambiguous lesion observed on CT scans (6).

Contrast-enhanced CT and MRI enable us to appreciate the anatomic relationship of the pulmonary veins and define the degree of anatomic stenosis by measuring vein size, allowing us to monitor and minimize any related complications (11). Post-acquisition processing from these modalities also allows 3D image reconstruction, which is beneficial for pre-procedural planning (5). In PVS, pulmonary parenchymal opacities and peripheral consolidations are typically indirect signs of significant stenosis and venous occlusion secondary to alveolar infarction or hemorrhage (15). Enhanced CT will reveal emboli as low-density structures within pulmonary arteries, and three-dimensional imaging will enable easier visualization at various angles. In our cases, no abnormal findings were observed in the lung parenchyma on the lung window CT scans.

Transcutaneous intervention for stenosis significantly improves the perfusion to the affected lung quadrant for most patients. However, normalization typically does not occur due to the varying degrees of irreversible injury sustained by the pulmonary vasculature

before stenosis relief (4). This was observed over time, in our case, using serial V/Q scans, which demonstrated partial and gradual improvement in perfusion after PTA. Based on CT findings alone, this improvement was not easily discernible in the first case.

Despite technological improvements and enhanced ablative techniques that have reduced the incidence of PVS from AF ablation, the morbid effects of a delayed diagnosis remain a clinical concern. Prompt diagnosis of PVS requires a high degree of suspicion due to its similarity to prevalent cardiopulmonary syndromes and the potential for misdiagnosis due to misleading diagnostic tests.

Studies have shown that using SPECT/CT has significantly improved the accuracy of lung perfusion scintigraphy (20). They are commonly applied for detecting pulmonary embolism and planning lung volume resection surgery for emphysematous and lung cancer patients. It addresses the previous limitations of V/Q planar and V/Q SPECT scans regarding anatomic and functional information. Lung perfusion scan combined with CT, performed by Hybrid SPECT/CT, has been found to offer a notably higher accuracy in detecting perfusion abnormalities compared to traditional scintigraphy techniques; however, it still lacks the necessary recommendations (20). Potential applications in PVS assessment may benefit PVS diagnostics. However, further investigation is still needed.

Conclusions

A V/Q mismatch in the form of reduced pulmonary perfusion is a notable finding of severe PVS and should be considered among patients with a history of atrial ablation. Along with CT, a V/Q scan is important in assessing PVS among patients with prior atrial ablation procedures. It can play a decisive role in characterizing the functional relevance of the stenosis and monitoring treatment response.

Acknowledgment

We thank The Otsuka Toshimi Scholarship Foundation for their support. JSPS KAKENHI, grant number 21K04080, partially supported this research.

References

1. Rostamian A, Narayan SM, Thomson L, Fishbein M, Siegel RJ. The incidence, diagnosis, and management of pulmonary vein stenosis as a complication of atrial fibrillation ablation. *J Interv Card Electrophysiol.* 2014; 40(1):63-74.

2. Prieto LR. The state of the art in pulmonary vein stenosis -diagnosis & treatment. *J Atr Fibrillation*. 2010; 2(4):228.
3. Raeisi-Giglou P, Wazni OM, Saliba WI, Barakat A, Tarakji KG, Rickard J, et al. Outcomes and management of patients with severe pulmonary vein stenosis from prior atrial fibrillation ablation. *Circulation: Arrhythmia and Electro-physiology*. 2018; 11(5): e006001.
4. Saad EB, Rossillo A, Saad CP, Martin DO, Bhargava M, Erciyes D, et al. Pulmonary vein stenosis after radiofrequency ablation of atrial fibrillation. *Circulation*. 2003; 108(25):3102-7.
5. Fender EA, Widmer RJ, Hodge DO, Packer DL, Holmes DR. Assessment and management of pulmonary vein occlusion after atrial fibrillation ablation. *JACC: Cardiovascular Interventions*. 2018; 11(16): 1633-9.
6. Nanthakumar K, Mountz JM, Plumb VJ, Epstein AE, Kay GN. Functional assessment of pulmonary vein stenosis using radionuclide ventilation/perfusion imaging. *Chest*. 2004; 126(2):645-51.
7. Simard T, Sarma D, Miranda WR, Jain CC, Anderson JH, Collins JD, et al. Pathogenesis, evaluation, and management of pulmonary vein stenosis: JACC review topic of the week. *Journal of the American College of Cardiology*. 2023; 81(24): 2361-73.
8. Mansour M, Gerstenfeld EP, Patel C, Natale A, Whang W, Cuoco FA, et al. Pulmonary vein narrowing after pulsed field versus thermal ablation. *EP Europace*. 2024; 26(2): euae038.
9. Lee JJ, Weinberg D, Anand R. Asymptomatic pulmonary vein stenosis: Hemodynamic adaptation and successful ablation. *Case Rep Cardiol*. 2016; 2016: 4979182.
10. Kumar N, Aksoy I, Pison L, Timmermans C, Maessen J, Crijns H. Management of pulmonary vein stenosis following catheter ablation of atrial fibrillation. *J Atr Fibrillation*. 2014; 7(1): 1060.
11. Pazos-López P. Pulmonary vein stenosis: Etiology, diagnosis and management. *WJC*. 2016; 8(1):81.
12. Bajc M, Schümichen C, Grüning T, Lindqvist A, Le Roux PY, Alatri A, et al. EANM guideline for ventilation/perfusion single-photon emission computed tomography (SPECT) for diagnosis of pulmonary embolism and beyond. *Eur J Nucl Med Mol Imaging*. 2019; 46(12): 2429-51.
13. Yun D, Jung JI, Oh YS, Youn HJ. Hemodynamic change in pulmonary vein stenosis after radiofrequency ablation: Assessment with magnetic resonance angiography. *Korean J Radiol*. 2012; 13(6): 816-9.
14. Holmes DR, Monahan KH, Packer D. Pulmonary vein stenosis complicating ablation for atrial fibrillation: Clinical spectrum and interventional considerations. *JACC: Cardiovascular Interventions*. 2009; 2(4):267-76.
15. Shroff N, Choi W, Villanueva-Meyer J, Palacio DM, Bhargava P. Pulmonary vein occlusion: A delayed complication following radiofrequency ablation for atrial fibrillation. *Radiol Case Rep*. 2021; 16(12): 3666-71.
16. Kind F, Goetz C, Meyer PT, Föll D, Russe MF, Ruf J. Pulmonary vein stenosis as a pitfall of ventilation/perfusion SPECT/CT for pulmonary embolism. *Clin Nucl Med*. 2021; 46(3): 238-9.
17. Yamaguchi K, Wakatsuki T, Matsuura T, Matsumoto K, Kawabata Y, Kadota M, et al. Drug-coated balloon angioplasty for severe pulmonary vein stenosis resulting from cryoballoon ablation for atrial fibrillation. *Journal of Cardiology Cases*. 2022; 26(1): 35-8.
18. Narechania S, Renapurkar R, Heresi GA. Mimickers of chronic thromboembolic pulmonary hypertension on imaging tests: a review. *Pulmonary Circulation*. 2020; 10(1): 2045894019882620.
19. Ostwani W, Arabi M. Alteration in pulmonary perfusion due to iatrogenic pulmonary vein stenosis: A mimicker of pulmonary embolism. *Avicenna J Med*. 2011; 1(2): 58-60.
20. Mazurek A, Dziuk M, Witkowska-Patena E, Piszczek S, Gizewska A. The utility of hybrid SPECT/CT lung perfusion scintigraphy in pulmonary embolism diagnosis. *Respiration*. 2015; 90(5): 393-401.