

# Importance of performing SPECT/CT in thyroid Cancer patients with central neck uptake in whole body Iodine scan: A case report

Nasrin Raeisi, Mohammad Ahmadi, Haniye Elahifard, Amin Saber Tanha\*

Nuclear Medicine Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

## ARTICLE INFO

*Article type:*

*Case Report*

*Article history:*

Received: 14 Sep 2025

Revised: 17 Nov 2025

Accepted: 21 Feb 2026

*Keywords:*

*Whole body iodine scan*

*SPECT/CT*

*Thyroid cancer*

*Pitfalls*

*Differentiated thyroid carcinoma*

## ABSTRACT

The planar whole body iodine scan (WBIS) has traditionally been used as the primary imaging technique for assessment of patients with thyroid cancer. This case report examines the imaging findings of a 61-year-old male with follicular thyroid carcinoma who underwent total thyroidectomy and received adjuvant radioactive iodine (RAI) treatment. A planar post-ablation WBIS revealed the commonly seen pattern of multiple foci of iodine uptake in the central neck region suggesting postsurgical thyroid remnant, thyroglossal duct remnant and central lymph node metastasis. However, single photon emission computed tomography (SPECT) / computed tomography (CT) imaging excluded lymph node metastasis and revealed iodine contamination over the mentum superimposed on the thyroid bed. This case report highlights the importance of utilizing SPECT /CT even for patients with apparently central neck uptake to prevent incorrect staging and treatment planning. In this report, we also present a review of the literature highlighting various pitfalls that can affect the interpretation of the WBIS and potentially result in false-positive findings in the neck region.

► Please cite this paper as:

Raeisi N, Ahmadi M, Elahifard H, Saber Tanha A. Importance of performing SPECT/CT in thyroid Cancer patients with central neck uptake in whole body Iodine scan: A case report. *Asia Ocean J Nucl Med Biol.* 2026; 14(2): .... doi:

## Introduction

Thyroid carcinoma is the most common endocrine malignancy and its incidence has been remarkably increased in the recent decades (1). The primary treatment for patients with differentiated thyroid carcinoma (DTC) is thyroidectomy followed by radioactive iodine (RAI) therapy in high risk patients (2). A post-ablation whole body iodine scan (WBIS) is mandatory for all patients undergoing RAI ablation and not only reveals local or distant metastasis, but also gives more information about the iodine avidity of the disease (3). The detection of lymph node or distant metastasis is the main tool for risk stratification. The extent and severity of the treatment highly depends on the risk of disease which is usually defined by TNM staging. The WBIS plays the main role for detection of local or distant metastasis, which is

traditionally performed as planar images. However, planar imaging has a number of limitations and single photon emission computed tomography (SPECT)/ computed tomography (CT) could potentially change TNM staging in a number of patients which could have therapeutic implications (4-6). In this case report we present a patient with suspicion of lymph node metastasis on planar scan which is excluded by SPECT/CT imaging and changed N category of TNM staging.

## Case presentation

A 61-year-old man with history of neck swelling for 4 months, underwent fine-needle aspiration (FNA) of a thyroid nodule in the right lobe that turned to be papillary thyroid carcinoma (PTC). Total thyroidectomy and central lymph node dissection was performed.


\* Corresponding author: Amin Saber Tanha. Nuclear Medicine Research Center, Nuclear Medicine Department, Qaem Hospital, Ahmadabad Avenue, Mashhad, Iran. Tel: +985138012779; Email: Saberta4011@mums.ac.ir

© 2026 *mums.ac.ir* All rights reserved.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The pathology examination showed a 3.5×2.4×2.2 cm follicular carcinoma in the right lobe with a few foci of poorly differentiated thyroid carcinoma. T and N category was pT2 and pN0a (none of the four resected central neck lymph nodes showed involvement). He was referred to our department for I-131 therapy. Upon presentation, the biochemical profile was as follows: serum thyroglobulin (S.TG): 2.59 ng/mL, serum anti-thyroglobulin (S. anti-TG): 20 IU/L, and thyroid-stimulating hormone (TSH) >78.5 mIU/L. Due to high risk features, he was admitted and 5.55 GBq of I-131 was administered as adjuvant therapy. A post-ablation planar WBIS was conducted seven days later utilizing a dual-head gamma camera (GE; Discovery NM/CT 670), equipped with high-energy collimators and configured with a 20% energy window set at 364 kiloelectron volts (keV). The scan was acquired in continuous acquisition mode at a table speed of 8 cm/min, employing automated body contour detection and recorded in a matrix size of 1024×256 (Figure 1-A). The planar scan revealed multiple foci of radio-iodine uptake in the central neck

region corresponding to the thyroglossal duct remnant (Figure 1-A, blue arrow), thyroid bed remnant (Figure 1-A, green arrow), and a focus inferior to thyroid bed, suggestive of lymph node metastasis (Figure 1-A, red arrow). SPECT imaging was performed from the neck region in a step-and-shoot mode (30 seconds/stop), 64 frames/head, using a noncircular orbit, over 360° (180° per head) saved in a 64×64 matrix and were reconstructed using iterative reconstruction (IR) method. CT acquisition with a tube voltage of 120 kV, tube current of 140 mAs and reconstructed slice width of 2.5 mm was performed subsequent to SPECT for attenuation correction, and anatomical localization. Contrary to findings of planar image, SPECT/CT images revealed that the uptake initially thought to be in the thyroid bed was actually an iodine contamination over the mentum. Furthermore, the focus previously misinterpreted as lymph node metastasis in planar WBIS, was confirmed to be the thyroid remnant on the left thyroid bed (Figure 1-B). So the patient was down staged from stage II (T2N1M0) to stage I (T2N0M0).



**Figure 1. (A)** The anterior projection of the whole-body radioiodine scan exhibits physiologic uptake patterns, accompanied by multiple sites of radioiodine accumulation in the central neck region, attributable to the thyroglossal duct remnant (**blue arrow**), iodine contamination on the mentum which was first considered as remnant in thyroid bed (**green arrow**), and residual thyroid tissue in the left thyroid bed which was assumed initially as possible lymph node metastasis (**red arrow**). **(B)** SPECT/CT imaging demonstrates that the uptake initially interpreted as thyroid bed remnant on the whole-body scan was, in fact, iodine contamination on the mentum (**green arrow**). Additionally, the focus regarded as possible lymph node metastasis on the whole-body iodine scan was confirmed to represent residual thyroid tissue in the left thyroid bed (**red arrow**). The remaining focus of radioiodine uptake was confirmed to represent the thyroglossal duct remnant, consistent with the initial presumption on the whole-body iodine scan (**blue arrow**).

## Discussion

Differentiated thyroid cancer, considered as the most prevalent endocrine malignancy, has increasingly become a significant public health concern. Globally, it ranked 7th most common cancer overall and 5th most common cancer in female, with projections indicating a potential increase of approximately 30 percent between 2019 and 2030 (7). Surgery remains the primary modality of treatment in these patients; however, internal radiation therapy utilizing RAI in high risk patients presents as important therapeutic necessity and the post-ablation WBIS is one of the main tools for risk stratification (8-12).

Traditionally WBIS is performed as planar imaging, however, false-positive findings and pitfalls are common and may affect risk stratification and treatment planning (13-18). Most of the patients with DTC have a small post-surgical thyroid remnant that can be seen as central neck uptake in a planar whole body iodine scan. Commonly, central neck activity is considered as thyroid remnant and/or thyroglossal duct remnant, and any extension beyond thyroid bed may suggest a lymph node metastasis (19). Additional views such as lateral

and/or oblique images may be helpful in occasional patients, however, due to poor spatial resolution of high-energy imaging, it is not easy to localize the lesion correctly in all of the patients. Although nuclear physicians are more prone to interpret central neck uptake as a thyroid bed remnant, false positive findings in this region might be seen and might result in incorrect diagnosis and treatment of the patient.

Numerous false positive iodine uptake has been reported in many organs either due to physiologic uptake or pathologic conditions. We summarized false positive uptake in the neck region in Table 1 (20-43). Our case report showed that saliva contamination may superimpose on the thyroid bed and be misinterpreted as a remnant or lymph node metastasis. Considering extensive cases of false positive in the neck region, SPECT/CT imaging could be invaluable for better delineation of any uptake in this region. This approach aligns with the recommendations in the latest edition of the American Thyroid Association (ATA) guidelines, which indicate that SPECT/CT may be performed when available in conjunction with diagnostic or post-treatment WBIS (44).

**Table 1.** Neck Iodine Activity (Review of the Literature)

<b>Thyroid-related uptakes</b>	<b>Physiologic</b>		<b>Pathologic</b>			
	1. <b>Ectopic thyroid</b>		1. Tumoral remnant in thyroid bed			
	a.	Pyramidal lobe (21,22)	2. Lymph node metastasis			
	b.	Aberrant tissue (21,22)	3. Cervical vertebrae metastasis(24)			
	c.	Lingual thyroid (21,22)	4. Cutaneous metastasis (25)			
	d.	Esophageal thyroid (23)				
	e.	Intra-tracheal thyroid (23)				
	2. <b>Post-surgical thyroid remnant (21)</b>					
	<b>Physiologic</b>					
	1. Salivary glands (21,26-28)					
2. Nasopharynx (26,28)						
3. Esophageal retention of iodine containing secretions (21,29)						
<b>Non-thyroidal uptakes</b>	<b>Pathologic</b>					
	<b>Infection/Inflammation</b>	<b>Benign Tumors</b>	<b>Malignant tumors</b>	<b>Trauma</b>	<b>Cystic degeneration</b>	<b>Other</b>
		Warthin's tumor (22,28,31)	Salivary gland adenocarcinoma (21,33)	Post traumatic superficial scab/wound (foreign body)(31,33)	Lymphoepithelial cyst of parotid gland (Sub-parotidian lymphoepithelial cyst) (22,33)	Barret's esophagous (28)
	Sialadenitis (22,30)	Salivary gland oncocytoma (27,31,34)	metastasis from other carcinoma (31)	Biopsy acupuncture sites (23,26,31)	Thyroglossal duct cyst (31,35)	Fistula tract (41)
	Dental disease (22,23,26,28)	Pleomorphic adenoma (35)	Squamous cell carcinoma (31)	Surgical suture(31)	Bone cysts (40)	Suture granuloma(41)
	Oral disease (22)	Spinal cord meningioma (33)		Surgical clips (28)		Laryngocele (21,42)
	Folliculitis (22,26,31)	Cavernous angioma (27,31)		Site of tracheotomy (31)		Sialolithiasis (27)
	TM joint effusion(32)	Vertebral hemangioma (31,36)		Post-operative seroma (38)		Salivary gland duct ectasia(27)
	Degenerative/inflammatory changes in Cervical vertebrae	Osteoid osteoma (31)		Bone fracture (39)		Dental amalgam (27)
	Skin burn(23,26,33)					
	Sebaceous cysts (23,29)	Lipoma (36)				Poorly dissolved <sup>131</sup> I capsule (21,23)
	Psoriatic plaque (23)	Nurilemoma (26)				Tracheostomy tube (26,28)
		Dermoid cyst (37)				Esophageal stricture (21,23,28)
						Zenker's diverticulum (21,23,28,31)
						Carotid ectasia (21-23)
	<b>External contamination*</b>	<b>Body secretions</b>				
		1. Saliva (22,23,26,43)				
		2. Nasal secretion (22,23,26)				
		3. Sweat (20,22,26,28,43)				
		4. Tracheobronchial secretions (22,26)				
5. Urine (23,26,28)						
6. Breast milk (23,26,28)						
7. Fecal (26)						
8. Vomitus (23,26)						

\* Signs of contamination: Handkerchief sign, Helmet sign, Necklace sign, Hot nose sign, Hot hand sign

In the present case, iodine contamination over the mentum, was superimposed on the thyroid bed and led to the misdiagnosis of lymph node metastasis in planar images. It was considered a TNM stage II disease, while SPECT/CT excluded

lymph node metastasis and downstaged it into TNM stage I. This case report underlines the importance of acquiring SPECT/CT images and using anatomic landmarks even in the common cases of central neck uptake in WBIS.

## Conclusion

This case highlights the essential role of SPECT/CT in distinguishing true central neck uptake from common pitfalls. Accurate localization in WBIS supports better risk stratification and treatment planning. We recommend SPECT/CT for all DTC patients if available, with careful consideration of potential false positives in the neck region.

## Acknowledgement

None.

## Conflict of interest

None to declare.

## Ethical consideration

The authors ensure that the work described has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans. The manuscript is in line with the Recommendations for the Conduct, Reporting, Editing and Publication of Scholarly Work in Medical Journals and aims for the inclusion of representative human populations (sex, age, and ethnicity) as per those recommendations.

## Contribution of author

NR & AS wrote the main text and gathered data. MA & HE provided the figures. AS revised the manuscript and is available for critical revision.

## References

- Huang J, Ngai CH, Deng Y, Pun CN, Lok V, Zhang L, et al. Incidence and mortality of thyroid cancer in 50 countries: a jointpoint regression analysis of global trends. *Endocrine*. 2023; 80(2): 355-65.
- Boucai L, Zafereo M, Cabanillas ME. Thyroid Cancer. *JAMA*. 2024; 331: 425.
- Haugen BR, Alexander EK, Bible KC, Doherty GM, Mandel SJ, Nikiforov YE, et al. 2015 American Thyroid Association management guidelines for adult patients with thyroid nodules and differentiated thyroid cancer: The American Thyroid Association guidelines task force on thyroid nodules and differentiated thyroid cancer. *Thyroid*. 2016; 26(1): 1-33.
- Nasr H, Alaklubi A, Mohamadien NR, Alqarni A, Farghaly H. The added value of SPECT/CT with radioactive iodine whole body scanning in patients with differentiated thyroid cancer and its impact on TNM staging. *Egyptian Journal Nuclear Medicine*. 2023; 26(1): 64-82.
- Mihailović J. Pre-treatment and post-treatment I-131 imaging in differentiated thyroid carcinoma. *Journal of Clinical Medicine*. 2024; 13(7): 1984.
- Holoubek SA, Sippel RS. Lymph node imaging for thyroid cancer. *Clinical Endocrinology*. 2024; 100(1): 96-101.
- Zhao Q, Chen M, Fu L, Yang Y, Zhan Y. Assessing and projecting the global burden of thyroid cancer, 1990–2030: Analysis of the Global Burden of Disease Study. *Journal of Global Health*. 2024; 14: 04090.
- Singh NK, Ramamourthy B, Hage N, Nagaraju S, Kappagantu KM. Radioactive iodine in differentiated carcinoma of thyroid: an overview. *Current Radiopharmaceuticals*. 2024; 17(1): 2-6.
- Chatterjee S, Mair M, Shaha AR, Paleri V, Sawhney S, Mishra A, et al. Current evidences in poorly differentiated thyroid carcinoma: a systematic review and subsection meta-analysis for clinical decision making. *Endocrine*. 2024; 85(2): 509-19.
- Jambi L, Aldawood Z, Alahmad H, Almuyid O, Qysi L, Alahmadi G, et al. The Role of Molecular Imaging in the Clinical Assessment of Well-Differentiated and Poorly Differentiated Thyroid Cancer. *Organization*. 2024; 1: 2.
- Berlińska A, Świątkowska-Stodulska R. Clinical use of thyroglobulin: not only thyroid cancer. *Endocrine*. 2024; 84(3): 786-99.
- Ran B, Shang J, Chen Y, Zhou M, Li H, He W, et al. The value of the first postoperative diagnostic I-131 scan in patients with papillary thyroid carcinoma. *Journal of Cancer Research and Clinical Oncology*. 2024; 150(2): 80.
- Buton L, Morel O, Gault P, Illouz F, Rodien P, Rohmer V. False-positive iodine-131 whole-body scan findings in patients with differentiated thyroid carcinoma: report of 11 cases and review of the literature. *In Annales d'endocrinologie* 2013; 74(3): 221-230.
- Agriantoni DJ, Hall L, Wilson MA. Pitfalls of I-131 whole body scan interpretation: bronchogenic cyst and mucinous cystadenoma. *Clinical Nuclear Medicine*. 2008; 33(5): 325-7.
- McDougall IR. Whole-body scintigraphy with radioiodine-131: a comprehensive list of false-positives with some examples. *Clinical Nuclear Medicine*. 1995; 20(10): 869-75.
- Shaik R, Hemalatha DS, Rallapeta RP, Sireesha P, Gavini ST, Kalawat T. Pitfalls of Iodine-131 Whole-Body Scan Mimicking

- Metastases in Differentiated Thyroid Carcinoma: A Case Series. *Indian Journal of Nuclear Medicine*. 2024; 39(1): 47-51.
17. Dillon M, Zielinski R, Worth J, Sanders M, Ibrahim O, Vedere T. An Unlikely Source of Iodine Uptake: A Bronchogenic Cyst Masquerading as Metastatic Thyroid Cancer. *JCEM Case Reports*. 2024; 2(3):luae042.
  18. Kamarulzaman K, Rohani MF, Nawi NM, Hassan SZ. Concurrent Benign Iodine-Avid Thymic and Liver Cysts Mimicking Metastatic Disease on <sup>131</sup>I Whole-Body Scintigraphy. *Clinical Nuclear Medicine*. 2024; 49(3): 250-2.
  19. Nava CF, Scheffel RS, Zanella AB, Zelmanovitz F, Maia AL, Dora JM. Reappraising the diagnostic accuracy of post-treatment whole-body scans for differentiated thyroid carcinoma. *Hormone and Metabolic Research*. 2020; 52(12): 834-40.
  20. Zakavi SR, Kakhki VD. Exercise-induced radio-iodine accumulation in scalp and hair during admission of <sup>131</sup>I therapy for thyroid cancer. *Thyroid*. 2006; 16(11):1185-6.
  21. Glazer DI, Brown RK, Wong KK, Savas H, Gross MD, Avram AM. SPECT/CT evaluation of unusual physiologic radioiodine biodistributions: pearls and pitfalls in image interpretation. *Radiographics*. 2013; 33(2): 397-418.
  22. Bakheet SM, Powe J, Larsson S, Hammami MM. Radioiodine uptake in the head and neck. *Endocrine Practice*. 2000; 6(1): 37-41.
  23. Shapiro B, Rufini V, Jarwan A, Geatti O, Kearfott KJ, Fig LM, et al. Artifacts, anatomical and physiological variants, and unrelated diseases that might cause false-positive whole-body <sup>131</sup>I scans in patients with thyroid cancer. In *Seminars in Nuclear Medicine*. 2000; 30(2):115-132.
  24. Ramadan S, Ugas MA, Berwick RJ, Notay M, Cho H, Jerjes W, et al. Spinal metastasis in thyroid cancer. *Head & neck oncology*. 2012; 4(1):39.
  25. McAninch EA, Calderon RM, Kargi AY. A cutaneous false positive in radioiodine scintigraphy for metastatic thyroid cancer. *US endocrinology*. 2016; 12: 37-8.
  26. Gholamrezaezhad A, editor. 12 Chapters on Nuclear Medicine. *BoD-Books on Demand*; 2011.
  27. Chudgar AV, Shah JC. Pictorial review of false-positive results on radioiodine scintigrams of patients with differentiated thyroid cancer. *Radiographics*. 2017; 37(1): 298-315.
  28. McDOUGALL IR. Whole-body scintigraphy with radioiodine-131: a comprehensive list of false-positives with some examples. *Clinical nuclear medicine*. 1995; 20(10): 869-75.
  29. Mitchell G, Pratt BE, Vini L, McCreedy VR, Harmer CL. False positive <sup>131</sup>I whole body scans in thyroid cancer. *The British Journal of Radiology*. 2000; 73(870): 627-35.
  30. Shapiro B, Rufini V, Jarwan A, Geatti O, Kearfott KJ, Fig LM, et al. Artifacts, anatomical and physiological variants, and unrelated diseases that might cause false-positive whole-body <sup>131</sup>I scans in patients with thyroid cancer. In *Seminars in Nuclear Medicine* 2000; 30(2): 115-132.
  31. Chambers MD, Khan MU. Interesting false positive radioiodine uptake on I-131 whole body scintigraphy with different mechanisms in two patients diagnosed differentiated thyroid carcinoma: A review of literature. *Clinical Reviews & Cases*. 2021; 3(1):1-5.
  32. Zhang M, Zhang Y, Huang W, Li B. False-positive <sup>131</sup>I uptake by the temporomandibular joint effusion. *Clinical Nuclear Medicine*. 2013; 38(10): 823-5.
  33. Buton L, Morel O, Gault P, Illouz F, Rodien P, Rohmer V. False-positive iodine-131 whole-body scan findings in patients with differentiated thyroid carcinoma: Report of 11 cases and review of the literature. In *Annales d'endocrinologie*. 2013; 74(3):221-230.
  34. Menon NN, Arora S, Dabas SK, Sharma A, Ranjan R, Gurung B, et al. Incidentally Detected <sup>131</sup>Iodine Avid Parotid Oncocytoma Coexistent with Papillary Carcinoma Thyroid. *Indian Journal of Otolaryngology and Head & Neck Surgery*. 2023; 75(3): 2598-603.
  35. Jia Q, Meng Z, Tan J, Zhang G, He Y, Sun H, et al. Retrospective imaging study on the diagnosis of pathological false positive iodine-131 scans in patients with thyroid cancer. *Experimental and Therapeutic Medicine*. 2015; 10(5): 1995-2001.
  36. Shaik R, Hemalatha DS, Rallapeta RP, Sireesha P, Gavini ST, Kalawat T. Pitfalls of Iodine-131 Whole-Body Scan Mimicking Metastases in Differentiated Thyroid Carcinoma: A Case Series. *Indian Journal of Nuclear Medicine*. 2024; 39(1): 47-51.
  37. Kosari HM, Zakavi SR, Barashki S, Firouzabad HR, Nakhaei SA, Aryana K. Incidental finding of a dermoid cyst in a whole-body iodine scan: importance of using [<sup>131</sup>I] SPECT/CT in the differentiated thyroid carcinoma. *Nuclear Medicine Review*. 2021; 24(2): 106-7.
  38. Aggarwal P, Seenivasagam RK, Sood A, Prashar S, Pathak P, Sachdeva N, et al. Radioactive iodine uptake in postoperative

- seroma: A cause for false positivity. *Journal of Nuclear Medicine Technology*. 2023; 51(1): 68-9.
39. Hannoush ZC, Palacios JD, Kuker RA, Casula S. False Positive Findings on I-131 WBS and SPECT/CT in Patients with History of Thyroid Cancer: Case Series. *Case reports in endocrinology*. 2017; 2017(1): 8568347.
40. Barbaro D, Campenni A, Forleo R, Lapi P. False-positive radioiodine uptake after radioiodine treatment in differentiated thyroid cancer. *Endocrine*. 2023; 81(1): 30-5.
41. Kandemir Z, Aksoy SY, Özdemir E, Aydın C, Yıldırım N, Türkölmez Ş. False positive result in the whole-body scan with 131-I in the follow-up of differentiated thyroid cancer: the fistula tract accompanied by the suture granuloma. *Gazi Medical Journal*. 2016; 27(3).
42. Schmidt M, Dietlein M, Schröder U, Schicha H. False-positive uptake of I-131 in a laryngocele mimicking thyroid remnant after thyroidectomy for papillary thyroid carcinoma. *Clinical nuclear medicine*. 2006; 31(11):716-7.
43. Anwariya A, Aggarwal P, Sood A, Tigapuram N, Prashar S. False positive finding on whole-body iodine-131 scan secondary to contaminated face mask: an uncommon peril in current COVID pandemic. *Asia Oceania Journal of Nuclear Medicine and Biology*. 2023; 11(1): 82.
44. Ringel MD, Sosa JA, Baloch Z, Bischoff L, Bloom G, Brent GA, et al. 2025 American Thyroid Association management guidelines for adult patients with differentiated thyroid cancer. *Thyroid®*. 2025; 35(8): 841-985.